

DOI: 10.14235/bas.galenos.2022.30602

Biological Activities and Chemical Composition of Turkish Sweetgum Balsam (*Styrax Liquidus*) Essential Oil
Türk Sığala Balzamu (*Styrax liquidus*) Uçucu Yağının Biyolojik Aktiviteleri ve Kimyasal Bileşimi

Büyükkılıç Altınbaşak et al. Biological Activities *Styrax Liquidus* Essential

Betül BÜYÜKKILIÇ ALTINBAŞAK¹, Ghassan ISSA², Belma ZENGİN KURT³, Betül DEMİRCİ⁴

¹Bezmialem Vakıf University, Faculty of Pharmacy, Department of Pharmaceutical Botany, Istanbul, Turkey

²Kocaeli Health and Technology University, Medical Laboratory Techniques Program, Avrupa Vocational School, Kocaeli, Turkey

³Bezmialem Vakıf University, Faculty of Pharmacy, Department of Pharmaceutical Chemistry, Istanbul, Turkey

⁴Anadolu University, Faculty of Pharmacy, Department of Pharmacognosy, Eskisehir, Turkey

Betül BÜYÜKKILIÇ ALTINBAŞAK, Bezmialem Vakıf University, Faculty of Pharmacy, Department of Pharmaceutical Botany, Istanbul, Turkey
betulbuyukkilic@gmail.com
0000-0002-9250-776X

17.12.2021

05.03.2022

Cite this article as: Büyükkılıç Altınbaşak B, Issa G, Zengin Kurt B, Demirci B. Biological Activities and Chemical Composition of Turkish Sweetgum Balsam (*Styrax Liquidus*) Essential Oil. *Bezmialem Science*.

ABSTRACT

Objective: The purpose of this present study was to make a chemical analysis of the composition of essential oil obtained from sweetgum balsam and to examine its antimicrobial, anticholinesterase, α -glucosidase inhibition, and antioxidant activities.

Methods: The essential oil obtained by the hydrodistillation method was analyzed by GC and GC/MS systems. The antimicrobial activities of the essential oil were evaluated by disc diffusion and resazurin microplate methods, the anticholinesterase effect on AChE and BChE enzymes, and the antioxidant effect by ABTS and CUPRAC methods.

Results: The main components of the essential oil were determined as styrene (92.6%) and α -pinene (2.2%). The essential oil showed weak antimicrobial activity against *K. pneumoniae*, *S. epidermidis* but strong antimicrobial activity against *A. baumannii*, *C. glabrata*. It showed

moderate inhibitory activity to AChE and BChE enzymes, and IC₅₀ values were calculated as 36.5 µg/mL and 69.5 µg/mL, respectively. It also showed low inhibition of α-glucosidase (IC₅₀ value 637.2 µg/mL) and a similar antioxidant effect in the CUPRAC and ABTS method (A_{0.5} value 637.2 µg/mL and IC₅₀ value 632.2 µg/mL, respectively).

Conclusion: Styrax Liquidus essential oil can be considered a natural antimicrobial agent due to its strong antimicrobial activity capacity against *A. baumannii* and *C. glabrata* strains.

Keywords: Styrax liquidus, essential oil, antimicrobial, anticholinesterase, α-glucosidase, antioxidant

ÖZ

Amaç: Bu çalışmada, sıgala balzamından elde edilen uçucu yağın bileşiminin kimyasal analizinin yapılması, antimikrobiyal, antikolinesteraz, α-glukozidaz inhibisyonu ve antioksidan aktivitelerinin incelenmesi amaçlanmıştır.

Yöntemler: Hidrodistilasyon yöntemiyle elde edilen uçucu yağ, GK ve GK/KS sistemleri ile analiz edilmiştir. Uçucu yağın antimikrobiyal aktiviteleri disk difüzyon ve resazurin mikropalak yöntemleri ile antikolinesteraz etkisi AChE ve BChE enzimleri üzerine, antioksidan etkisi ise ABTS ve CUPRAC yöntemleri ile değerlendirilmiştir.

Bulgular: Uçucu yağın ana bileşenleri stiren (%92.6) ve α-pinen (%2.2) olarak tespit edilmiştir. Uçucu yağ, *K. pneumoniae*, *S. epidermidis* kökenlerine karşı zayıf, *A. baumannii*, *C. glabrata* kökenlerine karşı güçlü antimikrobiyal aktivite göstermiştir. AChE ve BChE enzimlerine orta derecede inhibitör aktivite gösterip IC₅₀ değerleri sırasıyla 36,5 µg/mL ve 69,5 µg/mL olarak hesaplanmıştır. Ayrıca düşük oranda α-glukozidaz inhibisyonu (IC₅₀ değeri 637.2 µg/mL) ve CUPRAC ve ABTS yönteminde benzer bir antioksidan etki (sırasıyla A_{0.5} değeri 637.2 µg/mL ve IC₅₀ değeri 632.2 µg/mL) göstermiştir.

Sonuç: Styrax Liquidus uçucu yağı, *A. baumannii* ve *C. glabrata* suşlarına karşı güçlü antimikrobiyal aktivite kapasitesi nedeniyle doğal bir antimikrobiyal ajan olarak değerlendirilebilir.

Anahtar Sözcükler: Styrax liquidus, uçucu yağ, antimikrobiyal, antikolinesteraz, α-glukozidaz, antioksidan

Introduction

Styrax liquidus is a resinous exudate (sweetgum balsam) obtained from the wounded trunk of *Liquidambar orientalis* Mill. from the Altingiaceae family. The species is found only in southwestern Turkey and on the island of Rhodes. *L. orientalis* is an endangered relict species (1, 2). In history, Egyptian Queen Cleopatra's "love elixir" and used as perfume oil, essential oil of *Styrax liquidus*, has been used as a medicine since the Hippocrates. Ancient Egyptians used this oil during mummification. Amphoras filled with balsam extracted from sunken Phoenician ships indicate that the *Styrax liquidus* had an important place in the Mediterranean trade in the past (3).

The *Styrax liquidus* has been used for the treatment of various ailments in Turkish folk medicine. It is used to treat wounds, asthma, bronchitis, upper respiratory tract diseases, such as fumigation and an expectorant in the form of powder and pastille, skin diseases like scabies, and fungal in the form of pomade and plaster (4-7). The essential oil of *Styrax liquidus* has been used in the pharmaceutical and cosmetic industry particularly in perfumery. The essential oil constitutes 0.5-1% of the balsam (8).

In the literature review, it was determined that balsam has antibacterial, antioxidant and antiulcerogenic effects (9-11). *Styrax liquidus* essential oil has been reported to be inhibitory on the central nervous system as well as having an antimicrobial activity (12, 13).

The present study was undertaken to evaluate the antimicrobial, antioxidant, anticholinesterase, and α -glucosidase inhibition activities of *Styrax liquidus* essential oil and investigate its chemical composition.

UNCORRECTED PROOF

Methods

Extraction of essential oils

Styrax liquidus was obtained from local people in Marmaris, Muğla. The essential oil of *Styrax liquidus* was obtained by hydrodistillation in the Clevenger apparatus for 3 hours. Essential oil, thus obtained were stored in sealed vials at +4 °C until analyzed and tested.

Gas Chromatography (GC) and Gas chromatography-mass spectrometry analysis (GC/MS)

The composition of the essential oil was determined by using Gas Chromatography (GC) and Gas Chromatography/Mass Spectrometry (GC/MS).

GC analysis was performed on the Agilent 6890N GC system at an FID detector temperature of 300°C. To achieve the same elution order as GC/MS, the simultaneous automatic injection was performed on a replicate of the same column, applying the same operating conditions.

The relative percentage amounts of the separated compounds were calculated from the FID chromatograms and the analysis results are given in Table 1.

GC/MS analysis was performed on an Agilent 5975 GC/MSD system on an Innowax FSC column (60 m x 0.25 mm, 0.25 µm film thickness) using helium (0.8 ml/min) as carrier gas.

The GC oven temperature was held at 60°C for 10 minutes and programmed at 220°C at 4°C/min, and kept constant at 220°C for 10 minutes, then programmed to 240°C at 1°C/min.

The division ratio was set to 40:1 and the injector temperature was set to 250°C. Mass spectra were recorded at 70 eV. The mass range was m/z 35 to 450.

Identification of the essential oil components was carried out by comparison of their relative retention times with those of authentic samples or by comparison of their relative retention index (RRI) to a series of n-alkanes. Computer matching against commercial (Wiley GC/MS Library, MassFinder Software 4.0) (14, 15) and in-house “Başer Library of Essential Oil Constituents” built up by genuine compounds and components of known oils.

Antimicrobial activities

The essential oil was screened for their antimicrobial activities against *Streptococcus pneumoniae* ATCC 49619, *Enterococcus faecalis* ATCC 29212, *Staphylococcus aureus* ATCC 29213, *Staphylococcus. epidermidis* ATCC 49461, *Acinetobacter lwoffii* ATCC 19002, *Acinetobacter baumannii* ATCC 19606, *Klebsiella pneumoniae* ATCC 700603, *Pseudomonas aeruginosa* ATCC 27853, *Escherichia coli* ATCC 25922, *Candida albicans* ATCC 66027 and *Candida glabrata* ATCC 2001 microorganisms by disc diffusion and resazurin microplate methods.

Disk Diffusion Test

Fresh passages from the microorganisms were taken first to be tested using Disk Diffusion Tests in our study. The next day, suspensions were obtained from standard strains in saline water with a turbidity level of McFarland 0.5 (10⁸ microorganisms/mL). The Mueller Hinton Agar plate was seeded with a sample taken from these suspensions using a sterile swab. The essential oil of sweetgum balsam were first passed through a 0.22 µm filter for sterilization. Then, the paper discs obtained from Whatman paper that we would prepare were impregnated with the essential oil of sweetgum balsam placed on the plates. Moreover, various antibiotic discs with known sensitivity were also placed in Petri dishes for comparison. During this process, care was taken to keep a 22 mm space between the discs and a 14 mm distance from the edge of the Petri dish to ensure that the zones formed did not overlap. The inhibition zones were then measured by incubating the media for 18-24 hours at 37 °C.

Resazurin Microplate Assay (REMA)

Resazurin (7-hydroxy-3H-phenoxazin-3-one-10-oxide) microplate method was used to determine the antibacterial activities and minimum inhibitory concentrations (MIC) of the essential oil of sweetgum balsam *in vitro* against the standard origins of *Streptococcus*

pneumoniae ATCC 49619, *Enterococcus faecalis* ATCC 29212, *Staphylococcus aureus* ATCC 29213, *S. epidermidis* ATCC 49461, *Acinetobacter lwoffii* ATCC 19002, *Acinetobacter baumannii* ATCC 19606 *Klebsiella pneumoniae* ATCC 700603, *Pseudomonas aeruginosa* ATCC 27853 *Escherichia coli* ATCC 25922, *Candida albicans* ATCC 66027 and *C. glabrata* ATCC 2001. The activity determination was planned as two replicates. Streptomycin and fluconazole were used as the standard drugs. Stock solutions of the studied substances were prepared with DMSO at a concentration of 1024 µg/mL and sterilized by passing through a 0.22 µg membrane filter. To begin, each well was filled with 100 µL of Muller Hinton Broth medium. Serial dilutions of the prepared solutions were made by adding 1000 µg/mL to the first well of 96-well microplates (MIC range of chemicals 0.3–1000 µg/mL). The final concentration of the standard drug streptomycin was adjusted to 83 µg/mL and other standard drug fluconazole was adjusted to 30 µg/mL, and serial dilutions were made by adding 50 µL to the first well. Only DMSO was placed in one column of the plate as a negative control, and only standard bacteria as a positive control in the other column, both of which were 50 µL, and serial dilutions were made. A suspension of McFarland 0.5 turbidity was prepared from 1-2 day old colonies of microorganisms and then diluted 1:100. 10 µL of these final suspensions were added to the plate wells. Plates were covered with parafilm and incubated in a normal atmosphere for 24 hours at 37 °C. After the incubation, 10 µL each of 33.75 mg resazurin and 20% Tween 80, which were dissolved in 5 mL distilled water, were added to all wells, and plates were left to incubate for 2-4 hours, and the results were evaluated visually. The MIC value was determined to be the lowest concentration value preventing the color change from purple to pink.

Inhibitory activities against AChE and BChE

The anticholinesterase activity of essential oil was determined by using *in vitro* AChE and BChE enzymes inhibition assays. AChE and BChE inhibition activities were determined using the method found by Ellman et al. (16). Galantamine is used as reference compound. The IC₅₀ was determined by constructing an absorbance and/or inhibition (%) curve and examining the effect of seven different concentrations. Acetylthiocholine iodide and butyrylthiocholine iodide were used as substrates in the reaction, and 5,5'-dithio-bis(2-nitrobenzoic) acid (DTNB) was used as a reagent. Stock solutions of essential oils and galantamine were prepared with methanol at a 4000 µg/mL concentration. Acetylthiocholine iodide at 7.085 mM concentration and butyrylthiocholine iodide at 0.785 mM concentration were prepared. 150 µL of 100 mM phosphate buffer (pH 8.0), 10 µL of sample solution, and 20 µL of AChE (2.476x10⁻⁴ U/µL) (or 3.1813x10⁻⁴ U/µL of BChE) solution were mixed. It was incubated at 25°C for 15 minutes. 10 µL of DTNB solution at a concentration of 5mM was added. The reaction was initiated by the addition of 10 µL acetylthiocholine iodide (or butyrylthiocholine iodide). In this method, the activity was measured by following the yellow color produced as a result of the thio anion produced by reacting the enzymatic hydrolysis of the substrate with DTNB. Also, methanol was used as a control solvent. The hydrolysis of the substrates was monitored using a BioTek Power Wave XS at 412 nm (17). All experiments were done in triplicate and inhibition activity was calculated as;

Inhibition (%) = (Absorbance_{Control} - Absorbance_{Sample}) / Absorbance_{Control} x 100

IC₅₀ values were calculated using Graphpad Software.

α-Glucosidase inhibition assay

Commercially available α-glucosidase from *Saccharomyces cerevisiae* (Sigma, G5003) was selected as the target protein using p-nitrophenyl-α-D-glucopyranoside (pNGP, Sigma, N1377) as substrate. The essential oil and genistein were dissolved in DMSO, and the enzyme and substrate were dissolved in potassium phosphate buffer (0.05 M, pH 6.8). The enzymatic reaction mixture consisting of α-glucosidase (0.02 U, 20 µL), substrate (1.25 mM, 30 µL),

essential oil solution (10 μL), and potassium phosphate buffer (140 μL) was incubated at 37°C for 30 minutes. Next, the absorbance of the yellow color produced due to p-nitrophenol formation was measured at 405 nm using a Synergy H1 (BioTek, USA) 96-well microplate reader (18). All experiments were done in triplicate. α -glucosidase inhibition activity was calculated as;

$$\text{Inhibition (\%)} = (\text{Absorbance}_{\text{Control}} - \text{Absorbance}_{\text{Sample}}) / \text{Absorbance}_{\text{Control}} \times 100$$

IC₅₀ values were calculated using Graphpad Software.

Antioxidant activity assay

ABTS cation radical scavenging assay

ABTS cation radical scavenging activities of the essential oil were determined using 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulfonic acid) (19). A stock solution of 1000 $\mu\text{g}/\text{mL}$ of essential oil was prepared. 2, 5, 10, and 20 μL of this stock solution were taken and their volume was made up to 40 μL with ethanol, and 160 μL of 7 mM ABTS cation radical solution was added to them. After the reaction was kept in the dark for 6 minutes, absorbance was measured at 734 nm. ABTS cation radical scavenging activity was calculated by evaluating the absorbance values of the essential oil sample against the control. The percent inhibition was calculated from the following equation;

$$\text{Inhibition (\%)} = (\text{Absorbance}_{\text{Control}} - \text{Absorbance}_{\text{Sample}}) / \text{Absorbance}_{\text{Control}} \times 100$$

IC₅₀ values were calculated using Graphpad Software.

CUPRAC assay

In the presence of antioxidant compounds in the essential oil, Cu(II)-Neocuproin (Nc) complex was reduced to colored Cu(I)-Nc chelate, and the absorbance of this chelate was measured at 450nm. Butylated hydroxytoluene (BHT) was used as standard. CuCl₂, neocuproin, and NH₄OAc buffer were added to the essential oil and BHT with final concentrations of 10, 25, 50, 100 $\mu\text{g}/\text{mL}$, and absorbance were measured at 450 nm after 1 hour (20). The absorbance values of the essential oil sample were evaluated against the standard. The study was performed in three replications.

UNCORRECTED PROOF

Results

Composition of the essential oils

The analysis of essential oil obtained by the hydrodistillation method was carried out with GC/FD and GC/MS systems. Comparative analysis results are given in Table 1.

No	RRI	Compound	(%)
1	1032	α -Pinene	2.19
2	1035	α -Thujene	0.06
3	1076	Camphene	0.01
4	1118	β -Pinen	0.5
5	1203	Limonene	0.07
6	1218	β -Phellandrene	0.05
7	1272	Styrene	92.59
8	1466	α -cubebene	0.18
9	1497	α -Copaene	0.15
10	1541	Benzaldehyde	0.08
11	1549	β -cubebene	0.18
12	1611	Terpinen-4-ol	0.02
13	1612	β -caryophyllene	0.03
14	1671	Acetophenone	0.43
15	1704	γ -Murolen	tr
16	1740	α -Murolen	tr
17	1773	δ -Cadinene	0.19
18	1804	Myrtenol	tr
19	2049	4-Ethylguaiaicol	0.06
20	2065	Benzenepropanol	0.87
21	2068	(<i>E</i>)-Cinnamaldehyde	0.09
22	2113	Cumin alcohol	0.48
23	2157	(<i>E</i>)-Ethyl cinnamate	0.05
24	2195	4-Ethyl phenol	0.67
25	2219	α -Muurolol	0.24
26	2308	Cinnamyl alcohol	tr
Total		99.19	
Oil Yield		0.57	

RRI: Relative retention indices calculated against *n*-alkanes, % calculated from FID data, and tr: Trace (< 0.01%).

Antimicrobial activities

The essential oil was screened for their antimicrobial activities against *Streptococcus pneumoniae* ATCC 49619, *Enterococcus faecalis* ATCC 29212, *Staphylococcus aureus* ATCC 29213, *Staphylococcus epidermidis* ATCC 49461, *Acinetobacter lwoffii* ATCC 19002, *Acinetobacter baumannii* ATCC 19606, *Klebsiella pneumoniae* ATCC 700603, *Pseudomonas*

aeruginosa ATCC 27853, *Escherichia coli* ATCC 25922, *Candida albicans* ATCC 66027 and *Candida glabrata* ATCC 2001 microorganisms by disc diffusion method and resazurin microplate methods. According to our results; the essential oil showed weak antimicrobial activity against *K. pneumoniae*, and *S. epidermidis* but strong antimicrobial activity against *A. baumannii*, and *C. glabrata* (Table 2). The others microorganisms have not effectivite.

Table 2. Investigated microorganisms for antimicrobial effects

Microorganisms	Minimal inhibitory concentration of the essential oil of <i>Styrax liquidus</i> (($\mu\text{g/mL}$))
<i>Enterococcus faecalis</i> ATCC 29212	125
<i>Streptococcus pneumoniae</i> ATCC 49619	250
<i>Staphylococcus aureus</i> ATCC 29213	125
<i>Staphylococcus epidermidis</i> ATCC 49461	62.5
<i>Escherichia coli</i> ATCC 25922	125
<i>Pseudomonas aeruginosa</i> ATCC 27853	125
<i>Klebsiella pneumoniae</i> ATCC 70063	62.5
<i>Acinetobacter baumannii</i> ATCC 19606	31.25
<i>Acinetobacter lwoffii</i> ATCC 19002	125
<i>Candida albicans</i> ATCC 66027	125
<i>Candida glabrata</i> ATCC 2001	< 3.9

AChE and BChE inhibition

The essential oil moderately inhibited cholinesterase enzymes. Galantamine was used as the standard substance and the calculated IC_{50} values are given in Table 3.

Table 3. *In vitro* inhibition IC_{50} for AChE and BChE activities

Test sample	AChE IC_{50} ($\mu\text{g/mL}$)	BChE IC_{50} ($\mu\text{g/mL}$)
<i>Styrax liquidus</i> essential oil	36.5	69.5
Galantamine	0.55	3.56

α -Glucosidase inhibition

Styrax liquidus essential oil showed a low rate of α -glucosidase inhibition compared to genistein, which has strong inhibitory properties against α -glucosidase. Calculated IC₅₀ values are given in Table 4.

Test sample	IC ₅₀ (μ g/mL)
Styrax liquidus essential oil	637.2
Genistein	12.279

Antioxidant activity

Antioxidant effect was examined by CUPRAC and ABTS methods. Activity results were evaluated by comparison with standard BHT. It has been observed that essential oil has a very low antioxidant effect when compared to standard (Table 5).

Test sample	CUPRAC A _{0.5} (μ g/mL)	ABTS IC ₅₀ (μ g/mL)
Styrax liquidus essential oil	637.2	632.2
BHT	3.275	1.242
BHT: Butylated hydroxytoluene		

Discussion

Styrax Liquidus is known to have various biological effects due to its use in folk medicine and scientific studies. A limited number of studies have been reported the biological effects of essential oil Styrax liquidus. On GC-MS analysis of the essential oil, 26 compounds representing 99.19% of the total oil were identified where styrene (92.59%) and α -pinene (2.2%) were identified as the major components (Table 1).

There are several previous publications with different results in the literature regarding the chemical content of Styrax liquidus. Analysis by GC-MS resulted in cinnamyl cinnamate (21.5%), phenyl propyl cinnamate (7.5%), cinnamic acid (4.0%), cinnamyl alcohol (2.0%), styrene (0.5%) and phenyl propyl alcohol (0.5%). However, in this study, approximately 60% of the content was analyzed and different results are thought to be due to this (21). Another study reported that the main components of essential oil are styrene (89.5%), α -pinene (7.2%), and β -pinene (1.1%) (22). These results are very close to our findings.

In a different study, 58 components representing more than 99.4% of the essential oil and styrene (70.4%), α -pinene (19.4%), and β -pinene (4.3%) were determined as the main components (23).

The benefits of Styrax liquidus have been used for years, especially by local people, to treat skin problems, peptic ulcers, parasitic infections (4, 7, 24). The antibacterial, antiulcerogenic, antioxidant, and cytotoxic effects of styrax were investigated by researchers (10, 25, 26).

Our aim in this study is to reveal the potential for antioxidant, antidiabetic, anticholinesterase, and antimicrobial use of the essential oil of sweetgum balsam.

Anti- α -glucosidase compounds have received great attention due to their potential use in treating diabetes (27). This is the first report on anticholinesterase and α -glucosidase activities of essential oil of Styrax liquidus. The IC_{50} value against AChE and BChE is 36.5 μ g/mL and 69.5 μ g/mL, respectively. The IC_{50} value of enzyme inhibition results of galantamine is AChE=0.55 μ g/mL and BChE=3.56 μ g/mL, respectively. The obtained essential oil appears to inhibit cholinesterase enzymes at a moderate level. It showed a low rate of α -glucosidase inhibition (IC_{50} value of 637.2 μ g/mL, Genistein= IC_{50} value of 12.279 μ g/mL) and similar antioxidant effect on CUPRAC and ABTS method ($A_{0.5}$ value of 637.2 μ g/mL, BHT=3.275 μ g/mL, and IC_{50} value of 632.2 μ g/mL, BHT=1.242 μ g/mL, respectively).

A previous study investigated the antibacterial potential of Styrax liquidus against several bacteria strains by using the agar diffusion method. Different concentrations of balsam are potent on some bacterial strains. Its antibacterial activity in one of the most comprehensive studies; the concentration of 10% balm was effective against *Bacillus brevis*, *B. cereus*, *B. subtilis*, *Corynebacterium xerosis*, *Enterobacter aerogenes*, *Enterococcus faecalis*, *Klebsiella pneumoniae*, *Micrococcus luteus*, *Mycobacterium smegmatis*, *Proteus vulgaris*, *Pseudomonas aeruginosa*, *Pseudomonas fluorescent*, and *Staphylococcus aureus*. The concentration of 0.4% balm was effective against *E. aerogenes*, *P. vulgaris*. The concentration of 0.2% balm was effective against *E. aerogenes*, and *P. vulgaris*. Besides, the concentration of 0.1% does not show any antibacterial effect reported (11). In our study the essential oil of Styrax liquidus showed weak antimicrobial activity against *Klebsiella pneumoniae*, *Staphylococcus epidermidis* but strong antimicrobial activity against *Acinetobacter baumannii*, *Candida glabrata*. The others microorganisms have not effectivite (Table 2).

Due to their strong antimicrobial activities, plant-derived secondary metabolites are known to be important in the treatment of various diseases (28). In a study investigating the antiadhesive efficacy of coated with Styrax liquidus surgical silk sutures against common oral pathogenic microorganisms, the highest antiadhesive efficacy was observed against *S. aureus* (29).

Styrax liquidus is obtained from *L. orientalis*. This species is considered endangered due to threats such as agricultural activities, fires, pollution, polluted water, heavy tourism

investments, and overgrazing. It was therefore recorded as Critically Endangered on the 2017 European Red List (30).
In order to ensure continuity in the supply of natural products in pharmaceutical research, it is necessary to protect and cultivate medicinal plants.

UNCORRECTED PROOF

Study Limitations

Strong antimicrobial effects of *Styrax liquidus* essential oil have been determined. However, evaluation of its use as an antimicrobial agent has been limited, since studies on cytotoxic effects could not be performed.

UNCORRECTED PROOF

Conclusion

As in all balsams, the content of *Styrax liquidus* includes resin, essential oil, and free cinnamic acid. The resin containing "cytorezinol" constitutes 30-40% of the balm and 20-25% of the essential oil. The traditional medicinal use of the balm in various diseases and the low biological activity of the essential oil alone shows that the components in the balm have a synergistic effect. In this study, the most effective biological activity of essential oil is found as the antimicrobial effect. It is important to search for alternative antimicrobial agents because of resistance to existing antimicrobial drugs over time. *Styrax liquidus* essential oil can be considered a natural antimicrobial agent due to its strong antimicrobial activity against *A. baumannii* and *C. glabrata* strains.

UNCORRECTED PROOF

REFERENCES

1. Kavak S, Wilson B. *Liquidambar orientalis*. The IUCN Red List of Threatened Species 2018 [Internet]. <https://www.iucnredlist.org> [cited 02.02.2022]. Available from: <https://www.iucnredlist.org/species/62556/42326468>.
2. Öztürk M, Çelik A, Güvensen A, Hamzaoğlu E. Ecology of tertiary relict endemic *Liquidambar orientalis* Mill. forests. For. Ecol. Manag. 2008;256(4):510-8.
3. Deniz Ç. Comparison of “sigla oil”, a plant resin and collagenase containing pomades in wound healing (dissertation). Kahramanmaraş: Kahramanmaraş Sütçü İmam Unv. 2010.
4. Baytop T. Türkiyede bitkiler ile tedavi (geçmişte ve bugün) [Therapy with medicinal plants in Turkey, past and present]: Nobel Tıp Kitabevleri; 1999.
5. Ertuğ F. Bodrum Yöresinde Halk Tıbbında Yararlanılan Bitkiler. In: Başer KHC, Kıvrımer N. editors. BİHAT 2002. 14. Bitkisel İlaç Hammaddeleri Toplantısı; 2002 May 29-31; Eskişehir, Turkey; 2004. p.76-93.
6. Honda G, Yeşilada E, Tabata M, Sezik E, Fujita T, Takeda Y, et al. Traditional medicine in Turkey VI. Folk medicine in West Anatolia: Afyon, Kütahya, Denizli, Muğla, Aydın provinces. J. Ethnopharmacol. 1996;53(2):75-87.
7. Yeşilada E, Honda G, Sezik E, Tabata M, Goto K, Ikeshiro Y. Traditional medicine in Turkey IV. Folk medicine in the Mediterranean subdivision. J. Ethnopharmacol. 1993;39(1):31-8.
8. Tanker M, Sayron E. Styrax Liquidus Üzerinde Farmakognozik Araştırmalar. J. Fac. Pharm. Ankara 1974;(4):108.
9. Andrikopoulos NK, Kaliora AC, Assimopoulou AN, Papapeorgiou VP. Biological activity of some naturally occurring resins, gums and pigments against in vitro LDL oxidation. Phytother Res. 2003;17(5):501-7.
10. Gurbuz I, Yesilada E, Demirci B, Sezik E, Demirci F, Baser KH. Characterization of volatiles and anti-ulcerogenic effect of Turkish sweetgum balsam (*Styrax liquidus*). J. Ethnopharmacol. 2013;148(1):332-6.
11. Sağdıç O, Özkan G, Özcan M, Özçelik S. A Study on inhibitory effects of sığla tree (*Liquidambar orientalis* Mill. var. *orientalis*) storax against several bacteria. Phytother Res. 2005;19(6):549-51.
12. Koo B-S, Lee S-I, Ha J-H, Lee D-U. Inhibitory effects of the essential oil from SuHeXiang Wan on the central nervous system after inhalation. Biol. Pharm. Bull. 2004;27(4):515-9.
13. Topal U, Sasaki M, Goto M, Otles S. Chemical compositions and antioxidant properties of essential oils from nine species of Turkish plants obtained by supercritical carbon dioxide extraction and steam distillation. Int J Food Sci. 2008;59(7-8):619-34.
14. Hochmuth DH. MassFinder 4.0, Hochmuth Scientific Consulting. Hamburg, Germany 2008.
15. McLafferty FW, Stauffer DB. The Wiley/NBS Registry of Mass Spectral Data. New York: J. Wiley and Sons; 1989.
16. Ellman GL, Courtney KD, Andres Jr V, Featherstone RM. A new and rapid colorimetric determination of acetylcholinesterase activity. Biochem. Pharmacol. 1961;7(2):88-95.
17. Zengin Kurt B, Durdagi S, Celebi G, Ekhteiari Salmas R, Sonmez F. Synthesis, anticholinesterase activity and molecular modeling studies of novel carvacrol-substituted amide derivatives. J Biomol Struct Dyn. 2020;38(3):841-59. Epub 2019/03/07. doi: 10.1080/07391102.2019.1590243. PubMed PMID: 30836858.
18. Dege N, Özge Ö, Avcı D, Başoğlu A, Sönmez F, Yaman M, et al. Concentration effects on optical properties, DFT, crystal characterization and α -glucosidase activity studies: Novel Zn (II) complex. SAA. 2021;262:120072.

19. Sonmez F, Gunesli Z, Kurt BZ, Gazioglu I, Avcı D, Kucukislamoglu M. Synthesis, antioxidant activity and SAR study of novel spiro-isatin-based Schiff bases. *Mol. Divers.* 2019;23(4):829-44.
20. Apak R, Güçlü K, Özyürek M, Karademir SE. Novel total antioxidant capacity index for dietary polyphenols and vitamins C and E, using their cupric ion reducing capability in the presence of neocuproine: CUPRAC method. *J. Agric. Food Chem.* 2004;52(26):7970-81.
21. Hafizoglu H. Analytical studies on the balsam of *Liquidambar orientalis* Mill. by gas chromatography and mass spectrometry. *Holzforschung.* 1982;36(6):311-3.
22. Hafizoglu H, Reunanen M, Istek A. Chemical constituents of balsam from *Liquidambar orientalis*. *Holzforschung.* 1996;50(2):116-7.
23. Fernandez X, Lizzani-Cuvelier L, Loiseau AM, Perichet C, Delbecque C, Arnaudo JF. Chemical composition of the essential oils from Turkish and Honduras *Styrax*. *Flavour Fragr J.* 2005;20(1):70-3.
24. Lingbeck JM, O'Bryan CA, Martin EM, Adams JP, Crandall PG. Sweetgum: An ancient source of beneficial compounds with modern benefits. *Pharmacogn Rev.* 2015;9(17):1.
25. Atmaca H, Camli Pulat C, Cittan M. *Liquidambar orientalis* Mill. gum extract induces autophagy via PI3K/Akt/mTOR signaling pathway in prostate cancer cells. *International J. Environ. Health Res.* 2020:1-9.
26. Demir D, Özdemir S, Gonca S, Bölgen N. Novel *styrax liquidus* loaded chitosan/polyvinyl alcohol cryogels with antioxidant and antimicrobial properties. *J. Appl. Polym. Sci.* 2021:52033.
27. Athipornchai A, Kumpang R, Semsri S. Potential Biological Activities of *Clausena* Essential Oils for the Treatment of Diabetes. *J. Oleo Sci.* 2021;70(11):1669-76.
28. Burt S. Essential oils: their antibacterial properties and potential applications in foods-a review. *Int. J. Food Microbiol.* 2004;94(3):223-53.
29. Kiliç Y, Baygar T, Saraç N, Aysel U, Karaca İ. Anti-adhesion activity and physicochemical features of the surgical silk sutures coated with *Liquidambar orientalis* *styrax*. *Turkish Journal of Clinics and Laboratory.* 2020;11(5):359-65.
30. Rivers M, Beech E, Bazos I, Bogunić F, Buira A, Caković D, et al. European red list of trees: International Union for Conservation of Nature and Natural Resources (IUCN); 2019.