

Relationship Between Toilet Type and Hemorrhoids

Tuvalet Tipi ve Hemoroid Arasındaki İlişki

 Nedim UZUN¹,  Emine YILDIRIM²

¹Department of Emergency, University of Health Sciences Turkey, Gaziosmanpaşa Training and Research Hospital, Istanbul, Turkey

²Department of General Surgery, University of Health Sciences Turkey, Gaziosmanpaşa Training and Research Hospital, Istanbul, Turkey

ABSTRACT

Objective: The aim of the study is to investigate the relationship between the development of hemorrhoid disease and toilet habits.

Methods: This cross-sectional study was conducted at the digestive endoscopy unit of the Gaziosmanpaşa Training and Research Hospital between January 2022 and March 2022. Presence of constipation was evaluated according to Rome IV criteria and the patients were asked to fill out the short questionnaire after the procedure.

Results: There were a total of 142 patients in the study. The mean age was 53.6 ± 11.8 years. The female to male ratio was 1.33. The seated toilet has been reported by more than half of the patients (58.5%) as the preferred toilet type. Constipation was detected in 70 patients (49.3%). Hemorrhoidal disease was detected in 60 (42.3%) of the patients. We did not find a significant difference between hemorrhoid disease and the type of toilet in home, preferred toilet type and constipation ($p>0.05$). There were significantly more male patients with hemorrhoids (56.7% vs. 32.9%, $p=0.008$).

Conclusions: We concluded that there is no relationship between the preferred toilet type and hemorrhoid disease in the current study.

Keywords: Hemorrhoid, constipation, toilet type

ÖZ

Amaç: Çalışmanın amacı hemoroid hastalığının gelişimi ile tuvalet alışkanlıkları arasındaki ilişkiyi araştırmaktır.

Yöntem: Bu kesitsel çalışma Gaziosmanpaşa Eğitim ve Araştırma Hastanesi sindirim endoskopi ünitesinde Ocak 2022 ile Mart 2022 tarihleri arasında yapıldı. Konstipasyon varlığı Roma IV kriterlerine göre değerlendirildi ve işlem sonrasında hastalardan kısa anketi doldurmaları istendi.

Bulgular: Çalışmaya toplam 142 hasta alındı. Ortalama yaş 53.6 ± 11.8 yılı. Kadın erkek oranı 1.33 idi. Oturaklı tuvalet hastaların yarısından fazlası tarafından (%58,5) tercih edilen tuvalet tipi olarak bildirilmişti. 70 hastada (%49,3) kabızlık saptandı. Hastaların 60'ında (%42,3) hemoroidal hastalık saptandı. Hemoroidal hastalık ile evdeki tuvalet tipi, tercih edilen tuvalet tipi ve kabızlık arasında anlamlı bir fark saptanmadı ($p>0.05$). Hemoroidli erkek hasta sayısı anlamlı olarak daha fazlaydı (%56,7'ye karşı %32,9, $p=0.008$).

Sonuç: Bu çalışmada tercih edilen tuvalet tipi ile hemoroid hastalığı arasında bir ilişki olmadığı sonucuna vardık.

Anahtar Sözcükler: Hemoroid, kabızlık, tuvalet tipi

Introduction

Hemorrhoid disease is the third most common gastrointestinal system disease diagnosed in outpatient visits to the doctor in the USA (1). Hemorrhoids are diagnosed in one-third of cases undergoing colonoscopy (2). It is predicted that the demand for

hemorrhoid treatment will increase even more in the coming years (3).

Hemorrhoids are structures consisting of smooth muscle, connective tissue and vascular tissue that extend around the anal canal (4). Hemorrhoids are actually cushions that provide

Address for Correspondence: Emine YILDIRIM, Department of General Surgery, University of Health Sciences Turkey, Gaziosmanpaşa Training and Research Hospital, Istanbul, Turkey

E-mail: opdreylidirim@gmail.com **ORCID ID:** orcid.org/0000-0003-2733-402X

Received: 18.03.2022

Accepted: 30.06.2022

Cite this article as: Uzun N, Yıldırım E. Relationship Between Toilet Type and Hemorrhoids. Bezmialem Science

continence in healthy people (5). However, the concept of hemorrhoids is generally used for swollen blood bags and symptomatic hemorrhoid disease detected on examination, rather than a normal anatomical structure (4). The etiology of hemorrhoid disease is unclear (6). The most common view is that hemorrhoidal disease is caused by the hydrostatic pressure of the blood under the influence of gravity. The second opinion is that hemorrhoid disease occurs as a result of the reflection of arterial blood pressure in the anorectal region through the connections to the veins. Thirdly, it is thought that hemorrhoid disease occurs as a result of weakening of the anal mucosa due to recurrent irritation and inflammation (7).

Hemorrhoid disease is thought to be more common in developed countries, and it is assumed that this may be related to the low fiber diet and the use of seated toilets (8, 9, 10). It has been shown that the rectoanal angle is straightening in the squat posture (9), evacuation is easier and takes a shorter time (10). Turkey is a developing country between the west and east. Both seated and squat toilets are used. According to our hypothesis, we think that hemorrhoids may be significantly less in those who have the habit of using squat toilets.

Methods

The study was organized as a prospective cross-sectional study after local ethics committee approval.

Patients Selection

Consecutive outpatients who applied to Gaziosmanpasa Training and Research Hospital Endoscopy Unit for colonoscopy between January 2022 and March 2022 were included in the study. Hemorrhoids were diagnosed with colonoscopy by specialists in general surgery. Presence of constipation was evaluated according to Rome IV criteria (11). Toilet types at home were learned from all patients.

A questionnaire was applied to 161 patients whose age range was between 21 and 80 years. We excluded 19 patients with missing data and the presence of anorectal tumor, ileo-anal/rectal anastomosis, anal stenosis, anal fistula, anal fissure, pelvic floor disorders, any previous history of anorectal surgery, pelvic irradiation, neurologic diseases, diabetes, or chronic use of opioid medications., chron's disease, ulcerative colon and rectum tumors, as we thought that it would directly affect toilet habits. As a result, 142 patients were included in the study.

The relationship between the presence of hemorrhoids and age, gender, type of toilet in home, preferred toilet type and constipation were investigated.

Study Design

According to the pilot study results, which included a randomly selected ten patients from the database, 15.0% of this population has a hemorrhoidal disease. To estimate a 5% difference in effect size, with an error rate of 5% and 80% power, at least 53 patients are needed. Considering the possibility for incomplete or incorrect questionnaires by adding 10% loss to this figure results in a total of 58 patients needed.

Quantitative Variables

Etiologically, age, gender, fiber-poor diet, constipation, diarrhea and posture during defecation are thought to be associated with hemorrhoid disease (6). Cases with hemorrhoid disease were included in the research group, and cases without hemorrhoid disease were included in the control group.

Statistical Methods

Descriptive statistics were given as mean \pm standard deviation and median with minimum-maximum values for continuous variables depending on their distribution. Numbers and percentages were used for categorical variables. The normal distribution of the numerical variables was analyzed by the Shapiro-Wilk, Kolmogorov-Smirnov, and Anderson-Darling tests. The Independent Samples t-test was used to compare two independent groups where numerical variables had a normal distribution. Pearson Chi-Square and Fisher's Exact tests were used to comparing the differences between categorical variables in 2x2 tables. For statistical analysis, "Jamovi project (2021), Jamovi (Version 2.2.2.0) and JASP (Version 0.16) were used. In all statistical analyses, the significance level (p-value) was set at 0.05.

Results

There were a total of 142 patients in the study. The mean age was 53.6 ± 11.8 years. The female to male ratio was 1.33. The demographic and clinical characteristics of the study group are given in Table 1. The seated toilet has been reported by more than half of the patients (58.5%) as the preferred toilet type. Constipation was detected in 70 patients (49.3%). This rate was 43% in patients with hemorrhoids and 53.7% in patients without hemorrhoids. Other clinical characteristics of the patients are summarized in Table 1.

Of them, 60 patients (42.3%) reported the presence of hemorrhoidal pathologies. The mean age of the patients with and without hemorrhoids was similar ($p=0.654$). There were significantly more male patients with hemorrhoids (56.7% vs. 32.9%, $p=0.008$) Sex distribution and preferred toilet type of the patients with and without hemorrhoids shown in Figure 1 and 2. We detected no significant differences in the type of toilet present in their homes, preferred toilet type and constipation ($p>0.05$).

Discussion

Most studies in the literature on hemorrhoids are therapeutic, but studies on risk factors are few. There are studies showing relationship between age, gender, fiber diet, defecation posture, chronic constipation, portal hypertension, spinal cord injury and hemorrhoids (5). According to John F et al., the prevalence of hemorrhoids peaks between the ages of 45 and 65 and declines after the age of 65 (12). The effect of age and gender on anal pressure is known (13). Also, the relationship between anal pressure and hemorrhoids has been shown in many studies (14). In addition, anorectal mean resting pressure has been shown to

Table 1: Demographic and clinical characteristics of the study group

		Patients			p value
		Overall (n=142)	With hemorrhoids (n=60)	Without hemorrhoids (n=82)	
		mean± SS	mean± SS	mean± SS	
Age (year)		53.6 ± 11.8	54.1 ± 11.0	53.2 ± 12.4	0.654**
		n(%)	n(%)	n(%)	
Sex					
	Female	81 (57.0)	26 (43.3)	55 (67.1)	0.008*
	Male	61 (43.0)	34 (56.7)	27 (32.9)	
Type of toilet in home					
	Squat	71 (50.0)	30 (50.0)	41 (50.0)	0.999*
	Seated	107 (75.4)	41 (68.3)	66 (80.5)	0.143*
Preferred toilet type					
	Only squat+usually squat	55 (38.7)	26 (43.3)	29 (35.4)	0.154*
	Squat and seated	4 (2.8)	0 (0.0)	4 (4.9)	
	Only seated+usually seated	83 (58.5)	34 (56.7)	49 (59.8)	
Constipation					
	Positive	70 (49.3)	26 (43.3)	44 (53.7)	0.296*
	Negative	72 (50.7)	34 (56.7)	38 (46.3)	

*: Independent Samples T-Test, **: Pearson Chi-Square or Fisher's Exact test

decrease with age in 40 percent of studies (13). When we think logically, we expect the incidence of hemorrhoids to decrease with age. However, in the current study, no significant relationship was found between age and hemorrhoids. In women, on the other hand, it was found that the mean anal squeeze pressure was lower in most of the studies (15, 16). However, no significant relationship has been shown between external hemorrhoids and gender in adults (17, 18). But, it has been shown that boys have significantly more hemorrhoids (19). In our study, we found that hemorrhoids were significantly more common in men, as in boys.

There are few studies in the literature examining the relationship between toilet type and hemorrhoids, and the findings are controversial (9, 10, 19, 20). In the literature, we did not find a study similar to the current study in adults. It has been shown that the rectoanal angle is straightened during squat (9), in one study, defecation is more comfortable and less strained in the squat position, and in another study, complaints are reduced in hemorrhoid patients using the seated toilet(10). For this reason, it has been suggested that hemorrhoids can be seen less frequently in those who have squat toilet habits. According to a study by TuranYildiz et al., external hemorrhoids are significantly more common in children with seated toilet habits (19). But, in the current study, it was observed that there was no significant relationship between toilet type and hemorrhoid disease.

However, we should consider including both adults and patients with internal and external hemorrhoids in our study. Therefore, for only symptomatic hemorrhoid disease, a seated toilet may be a risk factor.

Chronic constipation is thought to be a risk factor for hemorrhoid disease (12). However, in our study, no significant relationship was found between the presence of constipation and hemorrhoids. In this study, it is thought that such a result was obtained because acute constipation was questioned, not chronic constipation. In addition, it would be more accurate to use scores such as KESS scoring in the evaluation of constipation (21).

As a result, studies on the etiology of hemorrhoids are insufficient, controversial and scarce in terms of evidence level. In the literature, the relationship between anal pressure and hemorrhoid disease has been revealed more clearly. Although we concluded that there is no relationship between the preferred toilet type and hemorrhoid disease in the current study, we think that this should be clarified by multicenter randomized cohort studies with long-term follow-up. It is thought that future studies on the causes of hemorrhoid disease will be designed by using the classification of hemorrhoids, BMI measurement, diet questioning, family history, anal manometry and chronic constipation scoring.




Figure 1: Sex distribution of the patients with and without hemorrhoids.




Figure 2: Preferred toilet types of the patients with and without hemorrhoids.

Study Limitations

Those included in the current study usually have a gastroenterological complaint, it would be better to have a similar study done on the general population. This is a cross-sectional study, it would be more appropriate to conduct a long-term cohort study. The grade and types of hemorrhoids were not included in the study, only the presence of hemorrhoids was examined.

Ethics Committee Approval: In this prospective study, ethics committee approval, was obtained from the Gaziosmanpaşa

Training and Research Hospital Ethics Committee, to which our hospital is affiliated (approval number: 49 and date: 22/12/2021).

Informed Consent: Informed consent was obtained from all patients participating in the study.

Peer-review: Externally and internally peer-reviewed.

Acknowledgement: I would like to offer my special thanks to Associate Professor Deniz Oğutmen Koc for their help and contribution to this paper.

References

1. Peery AF, Crockett SD, Murphy CC, Lund JL, Dellon ES, Williams JL et al. Burden and cost of gastrointestinal, liver, and pancreatic diseases in the United States: update 2018. *Gastroenterology*. 2019 May;156(6):1936. doi: 10.1053/j.gastro.2018.08.063
2. Everhart JE, Ruhl CE. Burden of digestive diseases in the United States part II: lower gastrointestinal diseases. *Gastroenterology*. 2009 Mar;136(3), 741-754. doi: 10.1053/j.gastro.2009.01.015
3. Etzioni DA, Beart RW Jr, Madoff RD, Ault GT. Impact of the aging population on the demand for colorectal procedures. *Dis Colon Rectum*. 2009 Apr;52(4):583-90. doi: 10.1007/DCR.0b013e3181a1d183.
4. Sun Z, Migaly J. Review of Hemorrhoid Disease: Presentation and Management. *Clin Colon Rectal Surg*. 2016 Mar;29(1):22-9. doi: 10.1055/s-0035-1568144.
5. Ganz RA. The evaluation and treatment of hemorrhoids: a guide for the gastroenterologist. *Clinical Gastroenterology and Hepatology* 2013 Jun; 11(6), 593-603. doi: 10.1016/j.cgh.2012.12.020
6. Sandler RS, Peery AF. Rethinking What We Know About Hemorrhoids. *Clin Gastroenterol Hepatol*. 2019 Jan;17(1):8-15. doi: 10.1016/j.cgh.2018.03.020.
7. Dimmer C, Martin B, Reeves N, Sullivan F. Squatting for the Prevention of Haemorrhoids?. *Townsend Letter for Doctors and Patients*. 1996. 159;66-71.
8. Burkitt DP. Hemorrhoids, varicose veins and deep vein thrombosis: epidemiologic features and suggested causative factors. *Can J Surg*. 1975 Sep;18(5):483-8.
9. Tagart RE. The anal canal and rectum: their varying relationship and its effect on anal continence. *Dis Colon Rectum*. 1966 Nov-Dec;9(6):449-52. doi: 10.1007/BF02617443.
10. Sikirov D. Comparison of straining during defecation in three positions: results and implications for human health. *Dig Dis Sci*. 2003 Jul;48(7):1201-5. doi: 10.1023/a:1024180319005.
11. Drossman DA. Functional Gastrointestinal Disorders: History, Pathophysiology, Clinical Features and Rome IV. *Gastroenterology*. 2016 Feb 19;S0016-5085(16)00223-7. doi: 10.1053/j.gastro.2016.02.032.
12. Johanson JF, Sonnenberg A. The prevalence of hemorrhoids and chronic constipation. An epidemiologic study. *Gastroenterology*. 1990 Feb;98(2):380-6. doi: 10.1016/0016-5085(90)90828-o.
13. Gundling F, Seidl H, Scalercio N, Schmidt T, Schepp W, Pehl C. Influence of gender and age on anorectal function: normal values from anorectal manometry in a large caucasian population. *Digestion*. 2010;81(4):207-13. doi: 10.1159/000258662.
14. Deutsch AA, Moshkovitz M, Nudelman I, Dinari G, Reiss R. Anal pressure measurements in the study of hemorrhoid etiology and their relation to treatment. *Dis Colon Rectum*. 1987 Nov;30(11):855-7. doi: 10.1007/BF02555423.
15. Sharma M, Lowry AC, Rao SS, Whitehead WE, Szarka LA, Hamilton FA, Bharucha AE. A multicenter study of anorectal pressures and rectal sensation measured with portable manometry in healthy women and men. *Neurogastroenterol Motil*. 2021 Jun;33(6):e14067. doi: 10.1111/nmo.14067.
16. Poos RJ, Bittner R, Frank J, Beger HG. Results of anorectal manometry for the determination of age- and sex-dependent pressure differences. *Zeitschrift für Gastroenterologie*, 1984 Oct,22(10), 592-597.
17. Greenspon J, Williams SB, Young HA, Orkin BA. Thrombosed external hemorrhoids: outcome after conservative or surgical management. *Dis Colon Rectum*. 2004 Sep;47(9):1493-8. doi: 10.1007/s10350-004-0607-y
18. Wronski K. Etiology of thrombosed external hemorrhoids. *Postepy Hig Med Dosw (Online)*. 2012 Jan 30;66:41-4.
19. Yildiz T, Aydin DB, Ilce Z, Yucak A, Karaaslan E. External hemorrhoidal disease in child and teenage: Clinical presentations and risk factors. *Pak J Med Sci*. 2019;35(3):696-700. doi: 10.12669/pjms.35.3.442.
20. Lam TC, Islam N, Lubowski DZ, King DW. Does squatting reduce pelvic floor descent during defaecation? *Aust N Z J Surg*. 1993 Mar;63(3):172-4. doi: 10.1111/j.1445-2197.1993.tb00512.x.
21. Knowles CH, Eccersley AJ, Scott SM, Walker SM, Reeves B, Lunniss PJ. Linear discriminant analysis of symptoms in patients with chronic constipation: validation of a new scoring system (KESS). *Dis Colon Rectum*. 2000 Oct;43(10):1419-26. doi: 10.1007/BF02236639.